

Figures

1.1	Interdisciplinary approach	8
2.1	Rural and urban monthly household incomes in Mongolia: 1997 to 2017	39
2.2	Per capita incomes in Inner Mongolia: 2000 to 2016	40
2.3	Livestock product and corn prices in Inner Mongolia: 2001 to 2016	42
3.1	Mean monthly air temperatures for Xilinhot, Inner Mongolia and Ulaanbaatar, Mongolia	51
3.2	Relationships between principal grassland species and average herbage mass over summer	55
3.3	Livestock numbers in Inner Mongolia and Mongolia over time	59
3.4	Relationship between stocking rates and animal production per head and per hectare	62
3.5	Energy balance for sheep equivalents on a typical farm in: (a) Siziwang <i>banner</i> , Inner Mongolia; and (b) Khashaat <i>soum</i> , Mongolia	65
3.6	Cumulative distribution functions for herder annual cash flows under different stocking rates: desert steppe, Inner Mongolia	70
3.7	Box plot of grassland biomass per hectare (summer and winter) under different stocking rates: desert steppe, Inner Mongolia	71
3.8	Box plot of meat production per hectare under different stocking rates: desert steppe, Inner Mongolia	72
3.9	Simulated dust emissions (winter and summer) under different stocking rates: desert steppe, Inner Mongolia	73
4.1	Radar plot of ratings for the steppe in Mongolia and typical steppe in Inner Mongolia	90
4.2	Direct primary links between the key eight grassland environmental services	92
4.3	Steps in grassland degradation identified in Mongolian grasslands and associated indicators	93
5.1	Four producer regions and three urban centres used in Mongolia price transmission analysis	98

5.2	Monthly Chinese and Mongolian beef and mutton prices (in CNY): January 2012 to August 2018	102
5.3	Indices of Chinese and Mongolian cashmere prices: 1998 to 2018	104
5.4	Monthly indices for mutton, beef and cashmere prices in China and Mongolia	106
6.1	Herder control beliefs: Mongolia	138
6.2	Herder power of control beliefs: Mongolia	139