1. Introduction

K.N. Ninan

ENVIRONMENTAL ASSESSMENTS: SCENARIOS, MODELLING AND POLICY

Environmental assessments are conducted to assess the environmental impacts of a policy or project and guide policy-makers and planners to assess alternative pathways to address, mitigate and internalize these adverse impacts. This could involve investing in a compensatory afforestation programme to compensate for the loss of a forest that is submerged or destroyed by a dam project, or installing an effluent treatment plant to treat the wastewater produced by an industry, or providing alternative livelihoods to local or indigenous communities who get displaced by an infrastructure project. A project may be financially viable but socially undesirable if the social costs exceed the social benefits. For instance, a hydro-electric project proposed in a tropical forest biodiversity hotspot may turn out to be financially viable if the social costs of the project are overlooked. But if the external costs of the project are accounted for, such as loss of forest area (including loss of biodiversity and ecosystem services) or indigenous communities displaced by the proposed project, the project may show negative returns. An environmental assessment will enable decision-makers to evaluate the trade-offs and how best to balance environmental concerns with development needs. The assessment could be an *ex ante* assessment where the appraisal is made prior to the likely implementation of the policy or project, or an *ex post* assessment where the appraisal is made after implementation of a policy or project.

There are a wide range of tools and frameworks to undertake environmental assessments. While environmental impact assessment (EIA) seeks to evaluate the likely environmental impacts at project level taking into account interrelated socio-economic, cultural and human health impacts, both beneficial and adverse, strategic environmental assessment (SEA) is broader in scope and makes an assessment at sectoral or geographical scales, with a systematic and comprehensive process of identifying and evaluating the environmental consequences of proposed

policies, plans or programmes to ensure that they are fully included and appropriately addressed at the earliest possible stage of decision-making on par with economic and social considerations (CBD, 2019). Integrated environmental assessment (IEA) includes environmental, social and economic aspects in an analysis of environmental state and trends linked with policy analysis. It usually covers a broad spectrum of issues and policies and all aspects of the environment including habitats, species and ecological, physical and chemical processes. It may incorporate global, sub-global and national perspectives as well as historical and future perspectives in an integrated analysis of environmental change and human and societal well-being (UNEP, 2019). Other assessment tools seek to assess the costs and benefits (including external costs and benefits) of a policy or project, such as social cost–benefit analysis (SCBA), multi-criteria analysis (MCA) or health and environmental risks, for example, risk assessment (RA) and comprehensive risk assessment (CRA) (Pearce et al., 2006). Tools such as environmental standards, emissions levels and ambient environmental standards are used to monitor and assess the impact of a project or programme on environmental quality such as air, water and soil quality (ADB, 2003). Other tools try to assess the contribution of the environment and natural assets to the economy, gross domestic product (GDP) and human well-being (for example, the United Nations' (UN) System of Environmental–Economic Accounting (SEEA), Green GDP, Genuine Progress Indicator (GPI)). A summary of selected environmental assessment tools and frameworks and their purpose is furnished in Table 1.1.

The focus of this book, however, is on scenarios and models and their usefulness for conducting environmental assessments and their role in decision-making and policy. Scenarios and models play different roles, but complement each other and fulfil a few important functions. They are useful in enlightening scientists, policy-makers and decision-makers on how the future is likely to unfold; alerting them about environmental risks such as global warming, loss of biodiversity and ecosystem services; helping to assess the alternative pathways to address these risks; helping in all phases of the policy cycle (agenda-setting and review, policy design and decision-making, policy implementation); and encouraging them to take proactive steps to address these risks, such as reducing use of fossil fuels, increasing the coverage of protected areas and restoring degraded ecosystems (IPBES, 2016). Since climate change has received considerable attention in scenario and modelling analyses for policy-making, our focus will be mainly on biodiversity and ecosystem services, which has received comparatively less attention.

Table 1.1 Summary of selected environmental assessment tools and frameworks

Environmental assessment tools and frameworks	Purpose
Environmental impact assessment (EIA)	To assess the environmental impacts of a proposed project or development, considering interrelated socio-economic, cultural and human health impacts, both beneficial and adverse; or a tool to identify the environmental, social and economic impacts of a project prior to decision-making (CBD, 2019).
Strategic environmental assessment (SEA)	Unlike EIA, SEA is broader in scope and can be applied to evaluating environmental impacts at sectoral or geographical scales. It seeks to identify and evaluate the environmental consequences of proposed policies, plans or programmes to ensure that they are fully included and appropriately addressed at the earliest possible stage of decision-making on a par with economic and social considerations (CBD, 2019).
Social cost–benefit analysis (SCBA)	SCBA tries to assess the social costs and benefits of a project including the income and distributional impacts of a project and its ability to fulfil development goals (e.g., employment generation, reduce poverty) as well as environmental costs and benefits. Unlike a financial analysis, which takes note of only financial costs and benefits, SCBA takes account of external costs and non-market benefits which need to be converted into monetary terms using economic valuation techniques.
Multi-criteria analysis (MCA)	MCA uses both quantitative and qualitative indicators to evaluate a project. While SCBA uses mono-criteria (such as net present value, benefit–cost ratio, internal rate of return) to appraise a project, MCA uses multiple criteria. For example, values attached to different aspects of biodiversity by stakeholders can be analysed using MCA to arrive at scores on different aspects of value. This gives a relative ranking to alternative notions of value such as use, bequest, option and existence values and provides a crucial input into stakeholders' perceptions with respect to biodiversity. However, a limitation of MCA is that when conflicting evaluation criteria are considered, an MCA problem may be mathematically ill-defined, making a complete axiomization and hence a simple decision criterion difficult to arrive at.

Table 1.1 (continued)

Environmental assessment tools and frameworks	Purpose
Life cycle assessment (LCA)	LCA is similar to EIA in that it identifies the environmental impacts of a policy or project and tries to measure them. The chief difference between EIA and LCA is that LCA looks not just at the impacts directly arising from a project or policy, but also at the whole 'life cycle' of impacts. For example, if the policy problem is to choose the best form of packaging a product, say fruit juice, the alternatives may be cartons, bottles and cans. LCA would look at the environmental impacts of each option, but going right back to the materials needed for manufacturing the container (e.g., timber, plastics, glass, metals) and the ways in which they will be disposed of once consumers consume the juice. LCA will include assessing the environmental impacts of primary resource extraction and the impacts from landfills (Pearce et al., 2006).
Risk assessment (RA)	RA involves assessing either the health or environmental risks (or both) attached to a product, process, policy or project (Pearce et al., 2006).
Comparative risk assessment (CRA)	CRA involves analysing risks for several alternative projects or policies and then suggesting opting for the one with lowest risk. For example, if the alternative is to choose between nuclear energy or coal-fired electricity, one approach is to normalize the risks of producing 1 kilowatt hour of electricity and to compute (probable) deaths per kWh. The option with the lowest 'death rate' may then be chosen. In case costs (say, cost of generating one kWh) are also to be considered, then cost-effectiveness analysis also needs to be conducted (Pearce et al., 2006).
Integrated environmental assessment (IEA)	An assessment that includes environmental, social and economic aspects in an analysis of environmental state and trends linked with policy analysis. It usually covers a broad spectrum of issues and policies, and all aspects of the environment including habitats, species and ecological, physical and chemical processes. It may incorporate global, sub-global and national

Table 1.1 (continued)

Environmental assessment tools and frameworks	Purpose
	perspectives, as well as historical and future perspectives, in an integrated analysis of environmental change and human and societal well-being (UNEP, 2019).
Environmental standards, emission levels, ambient environmental standards	To assess the predicted positive or negative changes brought about by project impacts on the environment. Environmental standards and emission levels are prescribed and monitored to assess the impact of an activity or project on environmental quality such as air, water and soil quality (ADB, 2003).
Green accounting, Green GDP	Green accounting seeks to make explicit the contribution of ecosystems and ecosystem services to economic activity. Also known as Green GDP; economic metrics that are ecologically adjusted by natural capital depletion can be produced by adjusting monetary measures of economic performance in response to impacts of changes to ecosystems such as the adjusted net national product (IPBES, 2016).
Inclusive or comprehensive wealth accounting	Involves estimating the changes in natural capital along with produced human and social capital. Future consumption possibilities such as ecosystem services depend on the various forms of capital and asset base of a nation (IPBES, 2016).
System of Environmental–Economic Accounting (SEEA)	A framework that integrates economic and environmental data to provide a more comprehensive and multi-purpose view of the interrelationships between the economy and the environment and the stocks and changes in stocks of environmental assets, as they bring benefits to humanity. It contains the internationally agreed standard concepts, definitions, classifications, accounting rules and tables for producing internationally comparable statistics and accounts. The SEEA framework follows a similar accounting structure as the System of National Accounts (SNA) (https://seea.un.org/).
Genuine progress indicator (GPI)	A metric designed to take fuller account of the well-being of a nation, only a part of which pertains to the health of

Table 1.1 (continued)

Environmental assessment tools and frameworks	Purpose
	the nation’s economy, by incorporating environmental and social factors which are not measured by gross domestic product (GDP). GPI has been suggested to replace or supplement GDP as a measure of economic growth (https://gnhusa.org/genuine-progress-indicator/).

Scenarios

Scenarios are often used to capture complexity, understand uncertainties, assess interactions of drivers of change, or to test alternative development trajectories (Priess and Hauck, 2014). There are varying definitions of ‘scenarios’, but on one point there is consensus: scenarios are not predictions of the future (van der Heijden et al., 2002). The Intergovernmental Panel on Climate Change (IPCC) describes scenarios as ‘a coherent, internally consistent and plausible description of a possible future state of the world. It is not a forecast; rather, each scenario is one alternative image of how the future can unfold’ (IPCC, 2014). The European Environment Agency (EEA) (2009) defines a scenario as a consistent and plausible picture of a possible future reality that informs the main issues of a policy debate. The methodological assessment report by the Intergovernmental Science–Policy Platform on Biodiversity and Ecosystem Services (IPBES) (2016), in the context of its mandate and focus on biodiversity and ecosystem services, describes scenarios as ‘representations of possible futures for one or more components of a system, particularly for drivers of change in nature and nature’s benefits, including alternative policy or management options’. Scenarios aim to foresee various pathways of changes in drivers of environmental change so that impacts on the environment or nature and nature’s benefits can be evaluated. Comparisons of alternative scenarios can then be used to guide and formulate policies such as increasing protected areas, reducing fossil fuel use, establishing hunting or fishing restrictions.

Scenario development emerged following World War II in US military strategic planning with the RAND Corporation. Scenarios achieved prominence in the 1970s in speculation about the future of society, the economy and the environment (van Notten, 2005). Today scenarios are used in a wide range of contexts by small, medium-sized and large

enterprises; in global, regional and national studies; and in environmental assessments for public policy, for example, the IPCC Assessment Reports, the United Nations Environment Programme's (UNEP) Global Environment Outlook, the Convention on Biological Diversity's (CBD) Global Biodiversity Outlook, the Organisation for Economic Co-operation and Development's (OECD) Environmental Outlook to 2050, the United Kingdom National Ecosystem Assessment (UK NEA) (Bateman et al., 2014; IPBES, 2016; van Notten, 2005).

Scenario development process

Scenario development involves building narratives or storylines of socio-economic and environmental pathways including assumptions regarding, for example, technological development, which can then be used to evaluate impacts on biodiversity and ecosystem services such as loss of biodiversity, species abundance, carbon sequestration, water availability, and so on (Pereira et al., 2010). These storylines are the backbones of the scenarios (Spangenberg et al., 2012). They are the qualitative component, defining philosophies, policies and instruments, which is then complemented by a quantitative component (Spangenberg et al., 2012). Sparrow (2000, cited in van Notten, 2005) argues that planners advising decision-makers should interpret scenarios as more exploratory, so that a scenario is less a strategy and more a coherently structured speculation. Of late, scenarios that are highly quantitative, including those based on econometric analyses, have gained prominence (Pereira et al., 2010).

The scenario development process involves consulting stakeholders, users and decision-makers; exploiting and improving the knowledge base (including indigenous and local knowledge) for developing scenarios; and evaluating their credibility and validity through peer reviews (IPBES, 2016). Once the contours of the scenarios are set and mapped, drivers are selected and interactions between drivers are analysed. The consequences of these social, economic and cultural developments, along with relevant data and other additional information, can be analysed by using models to project and estimate the values of the variables under study. Scenario analysis and models can enable governments and societies to anticipate future events (for example, global warming, loss of biodiversity and species) and the driving forces behind them (for example, fossil fuel use, habitat loss, land use change or pollution) and to take remedial measures (for example, reducing fossil fuel use, increasing protected area coverage, conservation programmes for species under threat of extinction) to address these environmental challenges (IPBES, 2016).

Types of scenarios

The scenario literature distinguishes between different types of scenarios based on their design and development process, such as exploratory and normative scenarios, qualitative and quantitative scenarios, deterministic and probabilistic scenarios, participatory scenarios, and so on (van Vuuren et al., 2012). Each approach has its advantages and limitations. For instance, in participatory approaches potential users work together with scenario developers so that participation can ensure that scenarios can be targeted better to user needs, and use available knowledge outside the group of directly involved experts (van Vuuren et al., 2012). However, participatory approaches may run the risk of losing more general applicability (van Vuuren et al., 2012). Explorative scenarios are meant to explore a wide range of possible futures, often to widen the scope of options considered by users (agenda-setting; strategy development); whereas normative scenarios focus on the impacts of implementing a more narrowly defined set of goals or policy options that often contribute to operational policy decisions (van Vuuren et al., 2012). Other approaches include the top-down versus bottom-up approach, where once scenarios are developed at a higher scale these are then used as a guide to develop the scenarios at lower scales (say, from global to regional and national scales); or vice versa, where after developing scenarios at lower scales these are aggregated to higher scales. The disadvantage of the first approach is that drivers or factors that may be relevant at, say, a global scale may not be that important at lower scales, and vice versa for the second approach. No approach can be considered as superior to another, and the choice of which approach to adopt for scenario development depends on the objectives of the scenario analysis and the needs and preferences of the users and decision-makers.

Considering the roles played by different types of scenarios in relation to the major phases of the policy cycle (agenda-setting, policy design, policy implementation and review) the methodological assessment report by the IPBES (2016) distinguishes between exploratory scenarios and target-seeking or policy-screening scenarios (Figure 1.1). While explanatory scenarios examine a range of plausible futures based on potential trajectories of drivers of environmental change that can contribute to high-level problem identification and agenda-setting, intervention scenarios evaluate alternative policy or management options through either target-seeking or policy-screening analysis, and can contribute significantly to policy design and implementation (IPBES, 2016). Exploratory scenarios also provide an important means of dealing with the high level of unpredictability, and therefore uncertainty, that is associated with the future trajectory of many drivers (IPBES, 2016). While exploratory scenarios have been widely used

Notes: Types of scenarios are illustrated by graphs of changes in nature and nature's benefits over time. The four phases of the policy cycle are indicated by arrows outside the circle. In 'exploratory scenarios' the dashed lines represent different plausible futures, often based on storylines. In 'target-seeking scenarios' (also known as 'normative scenarios') the diamond represents an agreed-upon target and the dashed lines indicate scenarios that provide alternative pathways for reaching the target. In 'policy-screening scenarios' (also known as 'ex ante scenarios') the dashed lines represent various policy options under consideration. In 'retrospective policy evaluation' (also known as 'ex post evaluation') the observed trajectory of policy implemented in the past (in solid black line) is compared to scenarios that would have achieved the intended target (dashed line).

Source: IPBES (2016, Figure SPM.2).

Figure 1.1 Roles played by different types of scenarios corresponding to major phases of the policy cycle

in assessments at the global, regional and national scales, intervention scenarios have been applied to decision-making mostly at national and local scales (IPBES, 2016).

Models

Models are a simplified depiction of complex economic or ecological processes. An economic model seeks to construct and analyse a mathematical model of the relationships and working of an economy or sub-sector of an economy. An ecological model (or ecosystem model) is the construction and analysis of mathematical models of ecological processes, including both purely biological and combined biophysical models (<https://www.nature.com/subjects/ecological-modelling>). They can be analytic or simulation-based, and used to understand complex ecological processes and predict how real ecosystems may change (ibid.). Models are constructed keeping in view logical consistency and theory, data availability, and objectives of the modelling such as to assess the factors that influence global temperature or sea-level or species abundance. Models can be static, that is, give a snapshot view of the economy or environment at a point of time; or dynamic, where time-related factors are also accounted for. Models are used to analyse and explain economic or ecological phenomena or used for prediction purposes.

The methodological assessment report (IPBES, 2016) defines models ‘as qualitative or quantitative representations of key components of a system and of relationships between these components’. In the assessment, ‘models’ usually, but not exclusively, refers to quantitative descriptions of relationships: (1) between indirect drivers and direct drivers; (2) between direct drivers and nature (including biodiversity and ecosystems); and (3) between nature and nature’s benefits to people (including ecosystem services) (IPBES, 2016).

Justification for using models or modelling

The justification for using models or modelling for undertaking environmental assessments are many. Environmental processes are complex, and hence need to be simplified and depicted through modelling which seeks to capture the relationships between different parameters of relevance (for example, between direct and indirect drivers of environmental change, or between nature, nature’s benefits and human well-being). Models are important as a scientific tool to support policy-making and decision-making. They can be used to assess and predict the impact of drivers on biodiversity and ecosystems, and hence the impacts on ecosystem services and human well-being. Furthermore, models (along with scenarios) can

help us to build plausible futures regarding the state of the environment, or alert us about future risks to the environment, biodiversity and ecosystem services, and help us to take informed decisions and assess various policy alternatives and take proactive measures (as against reactive measures) to address the risks posed by climate change, land use change and other drivers such as global warming, loss of biodiversity and ecosystem services. Models can also help us to assess past to present status, and trends in biodiversity and ecosystem services, and other parameters (IPBES, 2016).

Limitations and uncertainty

There are, however, a few limitations of models. One is the need to account for uncertainty – which is associated with all sciences, including modelling – due to data limitations, the representation of processes and the resolution of the ecosystem scale. The IPBES (2016) distinguishes between four categories of uncertainties, namely, ‘linguistic uncertainty’ (imprecise meaning, vagueness or ambiguity of words), ‘decision uncertainty’ (variations in subjective human judgements, preferences, beliefs and world-views), ‘stochastic uncertainty’ (random behaviour or unpredictability of complex natural, social and economic systems, particularly in relation to future states), and ‘scientific uncertainty’ (imperfect knowledge or data on the system being described). It is therefore quite common in IPCC and IPBES assessment reports to state the level of confidence attached to specific findings or statements, such as high, medium or low confidence, ‘well established’, ‘established but incomplete’, ‘inconclusive’, and so on. Another limitation is regarding the simplifying assumptions made to construct models to represent the real economy or environment. How realistic are they, and how far do they represent the real-world situation? Scaling issues, both spatial and temporal, are another issue of concern. For example, drivers have different impacts at different scales. The use of scenarios and models also varies, and depends on the different policy and decision contexts. No single model can capture all dynamics, and hence coupling of models is quite common. Biodiversity and ecosystem modelling depend heavily on our understanding of ecosystem structure, function and process, and on their adequate representation in models. Another issue is that current ecosystem services models do not address, or do not adequately address, the links or impacts of ecosystem services on human well-being and good quality of life (IPBES, 2016). Ecosystem services toolkits are weakly dynamic and do not incorporate ecological and social feedbacks (IPBES, 2016). For example, Integrated Model to Assess the Global Environment (IMAGE) and EcoPath with EcoSim (EwE) are useful for assessing current ecosystem services, but not well suited for addressing transformative change or long-term trends (IPBES, 2016).

Policy-makers have to make decisions even in the face of uncertainty, to act on drivers in order to conserve biodiversity and ecosystems. To support decision-making, models aim to synthesize this complexity into a reasonable number of dimensions. Sources of variability include natural variability (for example, genetic differences, and between environments), observation errors (for example, misspecified parameter distributions and biased parameter estimates) and structural complexity of a model due to many factors such as the number of parameters it includes, the number of ecosystem components and processes it simulates, the temporal scale; also non-linearities, threshold and cumulative effects incorporated in those effects, and so on (IPBES, 2016). These factors can increase ecosystem model uncertainty.

There are a number of ways to address uncertainty. One is to do a sensitivity analysis in order to test the robustness of the model parameters and estimated values. Another is to run a model multiple times to assess the full range of model behaviour, parameter uncertainty and natural variability. Lastly, it is suggested to use multi-model ensembles: for example, dynamic global vegetation model (DGVM) output could feed into species distribution models to better predict the reliance of species on biomes (IPBES, 2016).

Type of models and modelling approaches

I now review the various types of models and tools to assess biodiversity and ecosystem services. The IPBES methodological assessment report focused on models that address the effects of changes in indirect drivers (for example, socio-economic, political, technological and cultural factors) including policy interventions on direct drivers (for example, land use change, climate change, fishing pressure) of change in nature (biodiversity and ecosystem services); models that address the impacts of changes in direct drivers of nature; and those that address the consequences of changes in nature for the benefits that people derive from nature (IPBES, 2016).

Models to assess the impacts of drivers of change on biodiversity and ecosystem functioning can be classified based on biological or ecological levels (see Table 1.2). These models relate biological processes and environmental change to predict the existing and future state of biodiversity or ecosystem functioning at individual, species, community and ecosystem levels. Ecological interaction networks model species interactions such as predator–prey oscillations on a dynamic scale. Integrated assessment models (IAMs) rely on knowledge from diverse scientific disciplines to investigate and understand causal relationships within and between different sectors of the economy and the environment. Information about these model types and illustrations of their use are presented in Table 1.2.

Table 1.2 Models corresponding to different biological/ecological levels

Model type/biological or ecological level	Explanation	Illustrations of models used
Individual-level models and evolutionary adaptation	Theoretical models that can explicitly account for biological processes (e.g., mutation, dispersal and interactions within and between species such as mating and competition) and account for environmental change, and allow for projections of future biodiversity change scenarios beyond current conditions or observations.	For example, genetic variance models; evolutionary and population dynamic models.
Species or population-level models (also known as ecological niche modelling, habitat modelling, predictive habitat distribution modelling)	Biodiversity change at the species or population level is often measured using data on population demography and species distribution.	For example, logistic population modelling, life table matrix modelling, Lotka–Volterra models of community ecology, meta-population theory, equilibrium model of island biogeography, and so on, are the basis of ecological modelling to predict changes over time.
Community-level models	Enable to move beyond species-level predictions and predict broader impacts of environmental change. Like species distribution models they use environmental data to predict the distribution of species assemblages or communities.	Ferrier and Guisan (2006) identify three approaches to community-level distribution modelling: (1) assemble first, predict later; (2) predict first, assemble later; and (3) assemble and predict together (see IPBES, 2016, p. 136, Figure 4.4 for details).
Ecological interaction networks	Species interactions within communities can be explicitly modelled using process-based	Community interaction network approaches are used to assess impacts of invasive species,

Table 1.2 (continued)

Model type/biological or ecological level	Explanation	Illustrations of models used
	approaches that describe the links between species and the dynamics that determine species co-existence in the network, such as predator–prey oscillations.	overfishing of top predators, biodiversity and ecosystem function relationships, freshwater pollution, and so on. Species traits approaches are used to predict how changes in species or community traits will affect ecosystem functioning or assess ecological responses to natural and anthropogenic stressors based on species characteristics related to their functional role in ecosystems. Species–area relationships are used to predict impacts of changes in land use on species or community richness.
Ecosystem-level models, and integrated assessment models (IAMs)	Ecosystem-level models may focus on the biophysical dimensions of ecosystems or can be developed to include economic and social aspects. IAMs: an interdisciplinary process that combines, interprets and communicates knowledge from diverse scientific disciplines from the natural and social sciences to investigate and understand causal relationships within and between complicated systems (IPCC, 2001).	For example, dynamic global vegetation models (DGVMs) or EcoPath with EcoSim (EwE). For example, Integrated Model to Assess the Global Environment (IMAGE), Dynamic Integrated Model of Climate and the Economy (DICE). Shortcomings: may become too complex; largely focused on terrestrial systems and climate change.

Source: Brotons et al. (2016).

Models for assessing biodiversity and ecosystem services can also be classified in terms of process-based or correlative models, expert-based models and hybrid models (Table 1.3). Process-based models are those in which relationships are described in terms of explicitly stated processes or mechanisms based on established scientific understanding, and where model parameters have clear ecological interpretation, defined beforehand. In correlative models, available empirical data are used to estimate values for parameters that do not have predefined ecological meaning, and for which processes are implicit rather than explicit. Expert-based models are those in which the experience of experts and stakeholders, including local and indigenous knowledge-holders, is used to describe or model relationships. These models have their strengths and weaknesses. While process-based models represent a scientific understanding of key dynamics, which can enable learning and enrich decision assessment, their main weakness is that they require substantial knowledge and time to develop (IPBES, 2016). Correlative models are simple and easy to apply, but they are not based on process understanding and can dramatically fail in novel or data-poor situations (IPBES, 2016). Although expert-based models allow the relatively easy incorporation of diverse knowledge systems into ecosystem services models, expert knowledge is often partial, biased and can be incorrect, especially when applied to novel, complex or highly uncertain situations (IPBES, 2016). Hybrid models combine process and correlative models in order to represent complex, integrated systems with a focus on biophysical as well as human components. Some examples of the use of these model types are indicated in Table 1.3.

A summary of some major biodiversity models and modelling approaches and examples of their uses is presented in Table 1.4. These models differ in terms of the level of organization, typology, level of integration and expertise required, and the purposes for which they are used. These models relate environmental processes and factors such as age structure, population or area size, climatic and other factors to predict species occurrence, distribution and abundance, vegetation dynamics, water provision, carbon sequestration, crop production, and so on. While some of the models require a high level of expertise (for example, evolutionary models, DGVMs, IAMs), other models require basic or moderate level of expertise (for example, aquatic habitat suitability, species distribution models).

Some important models and tools to analyse ecosystem services are summarized in Table 1.5. While some tools are process-based (for example, IMAGE, EwE), others are correlative-based (for example, Co\$ting Nature, LUCI (Land Utilisation and Capability Indicator)), or expert-based (for example, ARIES (ARtificial Intelligence for Ecosystem Services), TESSA (Toolkit for Ecosystem Service Site-based Assessment)),

Table 1.3 Broad model types for modelling biodiversity and ecosystem services

Model types	Aims	Examples of use
Process-based models	Relationships are described in terms of explicitly stated processes or mechanisms based on established scientific understanding, and model parameters therefore have clear ecological interpretation, defined beforehand.	Hydrological process models are used to link land cover changes and land management to changes in quantity of freshwater supply and freshwater quality. Tree growth models with other parameters are used to simulate the carbon sequestration benefits of forests. Crop yield simulation models (e.g., Agricultural Production Systems Simulator, APSIM) use time series climate data, soil data and crop management regimes to simulate agricultural production in agro-ecosystems.
Correlative models	Available empirical data are used to estimate values for parameters that do not have predefined ecological meaning, and for which processes are implicit rather than explicit,	Assess the impacts of human activities on biodiversity, forecast future impacts of environmental changes, support human productive activities (e.g., enhance agricultural production) and conservation (Peterson, 2006; de Souza Muñoz et al., 2011).
Expert-based models	Experience of experts and stakeholders, including local and indigenous knowledge holders, is used to describe or model relationships.	Used in modelling species-habitat relationships, especially when data are scarce (e.g., Iglecia et al., 2012).
Hybrid models	Combine process and correlative models in order to represent complex, integrated systems with a focus on biophysical as well as human components (Parrott, 2011).	For example, the dynamic bioclimate envelope model developed by Cheung et al. (2008) simulates changes in the relative abundance of marine species through changes in population growth, mortality, larval dispersal and adult movement following the shifting of the bioclimate envelopes induced by changes of climatic variables. Climate envelope modelling is one model type that can be useful in understanding species and habitat responses to climate change because it identifies key links between drivers of change (e.g., climate) and relevant responses.

Source: Brotons et al. (2016).

Table 1.4 Summary of major biodiversity models and modelling approaches

Model	Level of organization	Model type	Level of integration	Required level of expertise	Examples of use	Reference
Evolutionary models	Organisms	Mixed (hybrid)	Integrated models	High	How demographic and evolutionary processes combine to predict whether a species will persist or not	Polechová and Barton (2015), Barton (2001)
Dynamic energy budget models	Organisms	Mechanistic	Integrated models	High	To understand evolution of metabolic organization	Kooijman (2009)
Aquatic habitat suitability	Species/ populations	Mainly correlative	Single or integrated models	Basic	To link environmental conditions to the quantitative habitat suitability of aquatic species	Mouton et al. (2009)
Species distribution models	Species/ populations	Mainly correlative	Single or correlative	Basic; moderate	Used to model the effects of environmental change on species distribution	Pearson and Dawson (2003), Elith and Leathwick (2009), Stockwell and Peters (1999), Phillips et al. (2006)
Community-level distribution models	Species	Mainly correlative	Integrated	Moderate	Use environmental data to predict species assemblages or communities (these models go beyond species predictions)	Ferrier and Guisan (2006)
Dynamic bioclimate envelope models	Species/ populations	Mixed (hybrid)	Integrated models	Moderate	Changes in the relative abundance of marine species induced by changes in climatic variables including population growth, mortality, larval dispersal, etc.	Cheung et al. (2008), Fernandes and Cook (2013), Gallego-Sala et al. (2010), Notaro et al. (2012)

Table 1.4 (continued)

Model	Level of organization	Model type	Level of integration	Required level of expertise	Examples of use	Reference
Age/stage structured models	Species/ populations	Correlative	Single	Basic	Widely used for fisheries management	Hilborn and Walters (1992), Getz (1988), Barfield et al. (2011)
Food web models	Ecosystems	Process- based	Integrated models	Moderate	Widely used for ecosystem-based management	Christensen and Walters (2011)
Size-based models	Community	Correlative models	Single model	Basic	In marine and freshwater ecosystems size is a good predictor of trophic level at community level	Duplisea et al. (2002), Rochet et al. (2011)
Species–area relationship models	Community	Correlative	Single models	Moderate	Used to predict the impacts of changes in habitat availability, driven by land use change or climate change on species; the most common species–area relationship model is the power function (Arrhenius, 1921), $S = cAz$, where S is species richness, A is habitat area, and c and z are model parameters (Rosenzweig, 2010)	van Vuuren et al. (2006), Desrochers and Kerr (2011), Pereira et al. (2013), Huth and Possingham (2011)
Biodiversity metric models	Community	Correlative	Integrated models	Moderate	A quantitative and integrated approach to assess the biodiversity with multiple indicators	Janse et al. (2015)

Lotka–Volterra	Community	Process-based	Integrated models	High	Used to examine interspecific competition and prey–predator interactions	Sinclair and Gosline (1997)
Dynamic global vegetation models (DGVMS)	Ecosystem	Process-based	Integrated models	High	To simulate the impact of climate change on vegetation dynamics at global scale and its carbon and water cycles	Botkin (1993), Bellard et al. (2012), Cramer et al. (2001)
General ecosystem models	Global	Process-based	Integrated models	High	Use a unified set of fundamental ecological concepts and processes for any ecosystem to which they are applied, either terrestrial or marine, at any spatial resolution	Harfoot et al. (2014)
Integrated assessment models (IAMs)	Global and regional	Integrated	Multiple models	High	Interdisciplinary assessment (presently used to model climate change and options for mitigation)	Harremoes and Turner (2001), Tilman et al. (2001)

Source: Brotons et al. (2016); for additional information on community-level distribution models, Simon Ferrier (personal communication).

Table 1.5 Summary of major ecosystem services models/tools

Tool	Model type	Scale in space, time	Ease of use	Community of practice	Flexibility	Reference
IMAGE	Process	Global, Dynamic	Difficult	Small	Low	Stehfest et al. (2014)
EcoPath With EcoSim (EwE)	Process	Region, Dynamic	Medium	Large	High	Christensen et al. (2005)
Artificial Intelligence for Ecosystem Services (ARIES)	Expert	Region, Dynamic	Difficult	Small	High	Villa et al. (2014)
Integrated Valuation of Ecosystem Services and Trade-offs (InVEST)	Process and correlative	Region, Static	Medium	Large	Medium	Sharp et al. (2018)
Co\$ting Nature	Correlative	Region, Static	Easy-Medium	Small	Medium	www.policysupport.org/costingnature
Toolkit for Ecosystem Service Site-based Assessment (TESSA)	Expert	Region, Static	Easy	Small	Low	Peh et al. (2014)
Corporate Ecosystem Services Review (ESR)	Expert	Region, Static	Easy	Small	Low	Hanson et al. (2012)
Land Utilisation and Capability Indicator (LUCI)	Correlative	Region, Static	Easy	Small	Medium	www.lucitools.org

Source: Peterson et al. (2016).

ESR (Corporate Ecosystem Services Review)). These tools also differ in terms of the scale of analysis (global or regional; dynamic or static). The table also throws light on the ease of their use, community of practice, and so on. For instance, InVEST, consisting of 18 software models developed by the natural capital project of Stanford University, is an open access free tool and is widely used to map and value ecosystem services. It helps to explore how changes in ecosystems can affect the flow of benefits to people. However, it is a data-hungry tool. More information about the uses of these models and tools is presented in Table 1.6. Table 1.7 throws light on the uses, strengths and weaknesses of a few expert-based models such as Bayesian belief networks, fuzzy cognitive maps, socioecological scenario analysis and matrix models to evaluate ecosystem services.

Policy

Environmental change driven by socio-economic and other factors is taking place at a rapid pace, and governments and societies need to anticipate these changes and respond effectively to address the likely adverse impacts of these changes such as rising global temperatures and sea-levels, growing frequency and intensity of disasters, and loss of biodiversity and ecosystem services. In this context scenarios and models can play an important role in enabling policy-makers and decision-makers to take informed decisions to address emerging environmental events and risks. Different types of scenarios play important roles in relation to the major phases of the policy cycle; that is, agenda-setting, policy design, policy implementation and policy review (IPBES, 2016). As stated earlier, while exploratory scenarios examine different plausible futures, based on potential trajectories of drivers, either indirect (for example, socio-political, economic and technological factors) or direct (for example, habitat conversion and climate change), and can contribute significantly to problem identification and agenda-setting, intervention scenarios that evaluate alternative policy or management options, through either ‘target-seeking’ or ‘policy-screening’ analysis, can contribute significantly to policy design and implementation. Models can provide a useful means for translating alternative scenarios of drivers of environmental change or policy interventions into projected consequences for nature (biodiversity and ecosystem services) and nature’s benefits to people (IPBES, 2016). The temporal and spatial scales at which scenarios and models need to be applied also vary across different policy and decision contexts (IPBES, 2016). Figure 1.2 gives an overview of the roles that scenarios and models play in informed policy and decision-making. Figure 1.3 depicts important characteristics and needs of policy and decision-making processes that are likely to vary

Table 1.6 Ecosystem services modelling tools and uses

Modelling type	Uses
InVEST (Integrated Valuation of Ecosystem Services and Trade-offs)	Widely applied suite of models for different ecosystem services, typically using the spatial extent and configuration of habitat or land use as predictors of ecosystem production in biophysical terms or economic terms; https://naturalcapitalproject.stanford.edu/software/invest
ARIES (Artificial Intelligence for Ecosystem Services)	A modelling platform incorporating multiscale process-based and probabilistic Bayesian models; it is spatially explicit, and any ecosystem service may be modelled.
EwE (EcoPath with EcoSim)	Used to dynamically represent energy flows through marine and aquatic systems; can assess only limited fisheries-related ecosystem services.
IMAGE 3.0 (Integrated Model to Assess the Global Environment)	IMAGE contains an ecosystem module that quantifies the supply of eight ecosystem services using other components in IMAGE 3.0 and also combine environmental variables and ecosystem services supply.
Co\$ting Nature	Used to estimate the cost of maintaining four ecosystem services (carbon storage, water yield, nature-based tourism and natural hazard mitigation).
ESR (Corporate Ecosystem Services Review)	Developed by the World Resources Institute (WRI); a structured methodology that helps businesses which interact with ecosystems to identify business risks and opportunities.
LUCI (Land Utilisation and Capability Indicator)	Uses simple algorithms and outputs to identify and communicate ecosystem services trade-offs to stakeholders and decision-makers.
TESSA (Toolkit for Ecosystem Service Site-based Assessment)	Uses decision trees to guide users through a process to rapidly prioritize ecosystem services for assessment and identify data needs and communication approaches.
PA-BAT (Protected Areas Benefits Assessment Tool)	A PDF/PowerPoint format tool to identify the range of ES and benefits. It uses a questionnaire approach in a workshop attended by different stakeholders/interest groups to identify PA Benefits. (Results based on local knowledge and not biophysical data. So may reflect biases of local stakeholders.)
MIMES (Multiscale Integrated Models of Ecosystem Services)	An analytical framework designed to integrate different ecological and economic models to understand and visualise ecosystem services (ES) values.

Table 1.6 (continued)

Modelling type	Uses
SolVES (Social Values for Ecosystem Services)	A geographical information system (GIS) based application for assessing, mapping and quantifying the social benefits of ecosystem services (ES), it derives a quantitative ten-point social value matrix from a combination of spatial and non-spatial responses to public value and preference surveys; it uses these data along with user-provided environmental data to map the provision of cultural ES across a region or landscape.
WaterWorld	A web-based spatially explicit model that comes with all the data required to run the model; focuses on water-related ES such as hydrological and water resources baseline and water risk factors with specific activities under current conditions and under scenarios for land use, land management and climate change.

Source: Neugarten et al. (2018).

markedly between contexts. The figure also depicts choices in the selection or design of scenarios, models and decision-support interfaces that depend on these policy context characteristics (IPBES, 2016). The use of scenarios and models for policy research and action, however, is constrained by gaps in data, knowledge, methods and tools (IPBES, 2016). Existing scenarios have some drawbacks, such as their failure to consider social-ecological feedbacks and multi-scale processes, and not explicitly addressing the links between targets for human development and for nature, and the interactions between global and local processes (Kok et al., 2017; Rosa et al., 2017). Efforts are under way to address these limitations of existing scenarios by placing nature at the centre of scenario development and addressing the full range of social-ecological feedbacks (Rosa et al., 2017).

ABOUT THE BOOK

Keeping in mind the above discussion, this book focuses on the theoretical and conceptual issues and applications of scenarios and models for assessing biodiversity and ecosystem services. The chapters included in this volume cover a broad range of topics such as scenarios, land use models, IAMs, participatory modelling, spatial models, climate models and their applications covering diverse ecosystems such as agriculture,

Table 1.7 Expert-based models

Model type	Uses	Strengths/weaknesses
Bayesian belief networks	Can be used to integrate expert knowledge with multiple data sources to model the flow of ecosystem services; they help to integrate expert and stakeholder knowledge with quantitative data and models.	Ability to combine different types of data and includes new data and its explicit treatment of stochastic uncertainty. Difficult to evaluate feedback processes; translating multiple types of data can be complex and confusing.
Fuzzy cognitive maps	These models are typically developed from discussions with experts, then iteratively revised into a model structure and function which corresponds to shared expert knowledge.	Can be used to make qualitative scenarios more rigorous and elicit models from diverse groups of people. Similar weaknesses as Bayesian belief networks.
Socio-ecological scenario analysis	A type of soft systems analysis to analyse the dynamics of social-ecological systems with a strong focus on ecosystem services and human well-being. Participatory scenario planning used to address social-ecological dynamics (participatory planning used to engage diverse communities to facilitate management and governance of landscapes for multiple benefits).	Unlike technical models, scenarios are more accessible, integrative and engaging; also, able to address trade-offs among different groups and multiple pathways between ecological change and human well-being. Less rigorous and comparable and less generalizable than technical models.
Matrix models	Used to integrate expert opinion, land cover data and other empirical data. For example, combining maps of land cover and land covers contribution to ecosystem services using GIS and matrices allows simple and rapid ecosystem services assessment.	Expert-based models allow relatively easy incorporation of diverse types of expert knowledge into ecosystem service models. Expert knowledge is often partial, biased and be incorrect especially when applied to novel, complex or highly uncertain situations.

Source: Peterson et al. (2016).

Notes: The left-hand panel illustrates how scenarios and models contribute to policy and decision-making through assessments, formal decision support tools and informal processes. Scenarios capture different policy options being considered by decision-makers which are then translated by models into consequences for nature, nature's benefits to people and quality of life. The left-hand panel also emphasizes that scenarios and models are directly dependent on data and knowledge for their construction and testing and provide added value by synthesizing and organizing knowledge (see bottom of left-hand panel). The right-hand panel provides an overview of the relationships between scenarios, models and key elements of the IPBES conceptual framework. The cross-sectoral integration elements signify that a comprehensive assessment of human well-being and good quality of life will often involve the integration of modelling from multiple sectors (for example, health, education, energy) addressing a broader range of values and objectives than those associated directly with nature and nature's benefits.

Source: IPBES (2016, Figure SPM.1).

Figure 1.2 An overview of the roles that scenarios and models play in informed policy-making and decision-making

Notes: Each arrow indicates that the selection or design of a particular attribute of ‘Assessment and decision-support interface’, ‘Scenarios’, ‘Models’ (right side of figure) is dependent on a particular characteristic or need of the ‘Policy or decision context’ (left side of figure).

Source: IPBES (2016, Figure 1.7).

Figure 1.3 Dependencies between the characteristics and needs of policy and decision-making processes in different contexts, and the selection or design of scenarios, models and decision-support interfaces to serve these needs

forests, wetlands, marine and urban areas. The chapters include those that have a global or regional focus as well as case studies drawn from Africa, Asia, Australia, Europe, North and South America. The contributors to this book are leading experts from around the world who have made a significant contribution to the literature in this area. Some of them have also been part of IPCC and IPBES assessments or other global, regional and national assessments. For convenience, the discussion in the book is organized into two parts: Part I: Theoretical and Conceptual Issues, and Part II: Applications and Case Studies. However, the discussion in some chapters in these two parts overlaps and covers both theoretical and conceptual issues and their application and uses.

Part I: Theoretical and Conceptual Issues

Canada is surrounded by three social-ecologically distinct ocean regions: the Arctic, Atlantic and Pacific. These oceans support rich biodiversity and provide vital social, economic and cultural benefits to Canadian society. The long-term sustainability of Canadian oceans is challenged by uncertainty over the impacts of future climate and socio-economic change. Scenario analysis can be used to address this uncertainty by exploring alternative futures for Canada's three oceans under different pathways of climate, economic development, social and policy changes. However, to date there has been no national-scale scenario developed to enable the investigation of Canada's future ocean sustainability. To facilitate this process, in Chapter 2, Teh et al. discuss whether existing scenarios of Canadian oceans provide an integrative, social-ecological perspective on potential future conditions for Canada's fisheries and marine ecosystems. They then apply the findings to inform the development of a national-level scenario framework which allows a social-ecological examination of Canada's oceans in terms of the state of future environmental, social and economic uncertainties.

IAMs are cross-disciplinary tools that explore how economic activity interacts with the environment and vice versa. They differ in their design and scope and are frequently used by the IPCC, national environmental agencies and others to assess the economic consequences of climate change. One group of IAMs goes as far as estimating the total economic effect of climate change, which includes three essential components: costs of emissions abatement, costs of adaptation to changing climate and residual impacts on the economy. The most prominent models in this group – Dynamic Integrated Model of Climate and the Economy (DICE), the Climate Framework for Uncertainty, Negotiation and Distribution (FUND) and Policy Analysis of the Greenhouse Effect (PAGE) – follow a

commonly adopted approach to use empirical damage functions as a way of translating climate-driven environmental changes into the economic impacts, which could be either costs or benefits. Chapter 3 by Prieg and Yumashev explores how damage functions are constructed and calibrated in these three IAMs by examining model documentation, and summarizes the limitations of the current approaches. The chapter then reflects on how damage functions could be improved in order to help build confidence in these IAMs, and ultimately provide better estimates of the impacts of climate change for decision-makers and academics.

Land use and land cover change (LULCC) is now one of the major drivers of biodiversity change, contributes significantly to climate change and ultimately impacts upon human well-being. Therefore, it is crucial to understand how land use change evolved over time and to project expected trajectories of future change, so that preventive and/or adaptive actions can be taken by decision-makers and policy-makers. In Chapter 4, Isabel Rosa first reviews the literature to highlight major advancements in understanding historical trajectories of LULCC across the globe. Then she discusses how different methodologies (for example, remote sensing, citizen science) have allowed us to greatly improve our ability to monitor present LULCC, drawing attention to current limitations and emerging solutions. Finally, she presents and compares several approaches (for example, IAMs, statistical models) to project scenarios of future LULCC at multiple spatial scales (from local to global). Further, Rosa discusses the ways forward to produce better LULCC projections that are policy-relevant and can be used as tools to help support decisions aiming at sustainable management of the world's natural resources. The chapter will enable the reader to get a holistic overview of the state of the art in the field of LULCC, from detection (monitoring) to scenario projections (modelling).

Part II: Applications and Case Studies

Ida Kubiszewski et al. in Chapter 5 estimate the future value of ecosystem services in monetary units for four alternative global land use and management scenarios based on the Great Transition Initiative (GTI) scenarios to the year 2050. The authors used previous estimates of the per biome values of ecosystem services in 2011 as the basis for comparison. They then mapped projected land use for 16 biomes at 1 km² resolution globally for each scenario. This, combined with differences in land management for each scenario, created estimates of global ecosystem services values that also allowed for examinations of individual countries. Results show that under different scenarios the global value of ecosystem services can

decline by US\$51 trillion/yr or increase by US\$30 trillion/yr. In addition to the global values, the report totals for all countries and maps for a few example countries. Results show that adopting a set of policies similar to those required to achieve the UN Sustainable Development Goals (SDGs) would greatly enhance ecosystem services, human well-being and sustainability.

The environmental value of protected areas is seriously threatened by the invasion of exotic weeds, particularly in countries such as Australia. The value of these natural assets for biodiversity conservation, and for recreation and tourism, is reduced by the presence of weeds in these areas, and ecosystem services and functioning are undermined. Resources for managing these weeds are limited, and therefore it is important to establish priorities for their control. Chapter 6 by Tisdell et al. outlines economic and ecological models that may be (or have been) used for determining these priorities, and assesses their limitations. The scope for applying SCBA and cost-effective analysis is explored, as well as less ambitious economic models. The use of ecological modelling based on the extent to which an invasive weed becomes established in protected areas is given particular consideration. This modelling is based upon the relevance of sigmoidal invasion curves. As pointed out, sometimes the implications of economic models for establishing priorities for weed control are at odds with policy conclusions drawn from ecological models. Actual processes for prioritizing weed control in some of Australia's national parks are investigated. It is found that pressure groups can have a substantial influence on the determination of these priorities.

Nationally determined contributions (NDCs) are central to the Paris Agreement on climate change, allowing Annex II (largely developing) countries to quantify voluntary contributions to the global mitigation effort. Research has been quick to identify the headline NDC figures: if these climate action plans were fully implemented, global mean warming by 2100 would be reduced from approximately 3.6°C to 2.7°C above pre-industrial levels. However, beyond these headline mitigation figures, many NDCs are difficult to analyse, and few include a contribution from the significant sector of agriculture, forestry and land use change. Brazil is the first developing country to provide an absolute emission cut as its NDC, seeking to reduce greenhouse gas emissions by 37 per cent below 2005 levels by 2025, and 43 per cent by 2030. The Brazilian NDC is also noteworthy in focusing on emissions from deforestation control and land use change and agricultural intensification. Chapter 7 by Silva et al. outlines the optimization modelling to develop mitigation scenarios that underpinned Brazil's NDC commitment.

Tropical countries such as Cambodia require information about forest

cover and biomass for successful implementation of climate change mitigation policies related to Reducing Emissions from Deforestation and Forest Degradation (REDD) as well as implementation of various SDGs. In Chapter 8, Avtar et al. examine deforestation in Cambodia and develop a spatial model to simulate future forest cover under a business-as-usual (BAU) scenario. Spatial modelling tools have been used to model present conditions and future scenarios. The results showed that the current deforestation process is in a critical stage where some sub-regions may face unprecedented stress on forests. Phased Array type L-band Synthetic Aperture Radar (PALSAR) data were also used to estimate above-ground biomass (AGB) in Cambodia for 2009 and 2015. In this research, Cambodia's forest cover from 2020 to 2040 was forecasted, and forest biomass for 2009 and 2015 was estimated. This forecasted forest cover and biomass information will be useful to establish forest reference emission levels (RELs) and/or forest reference levels (RLs) in Cambodia.

Changes in ocean physics and chemistry under climate change are already impacting upon marine organisms, ecosystems and the services they provide us, including seafood. This impacts upon the waters of British Columbia (BC), Canada, and will have direct impacts on supply of the staple seafoods that are consumed by residents of BC. Sumaila and Lam in Chapter 9 discuss how these changes would affect seafood household budgets of British Columbians under climate change. The study projects that prices of iconic west coast species will increase under a climate change scenario. The net change in price attributable to climate change costs British Columbians up to \$110 million a year in 2015 dollars, and this imposes a significant burden on household budgets. The impacts of climate change on the oceans (including temperature increase, acidification and deoxygenation) provide compelling arguments for rapid reductions in carbon dioxide (CO₂) emissions. To effectively mitigate the CO₂ emissions requires immediate actions and collaborations among different levels, from individual and private actors to the government and the intergovernmental levels.

The environmental conditions in Latin America and the Caribbean (LAC) have considerably eroded since 1960 as consequence of a combination of multiple factors and drivers. Chapter 10 by Ramón Pichs-Madruga mainly focuses on a sub-set of the socio-economic (indirect) drivers since 1960, and discusses some expected trends for the next decades. The assessment of these drivers is critical in providing inputs in terms of data and narratives for an environmental scenario-building process, as a key tool for policy-makers. The main regional priorities for LAC regarding the linkages between environment and development refer to the need for transformational changes in the regional socio-

economic agenda with a long-term perspective. In the absence of such transformational changes in the prevailing production and consumption patterns in LAC, most of the adverse trends and imbalances in socio-economic development, and their interlinkages with environmental problems, would intensify towards 2050 and beyond. This situation would compromise the regional progress towards Agenda 2030 and the accomplishment of the SDGs.

Passively crowdsourced social media act as an opportunistic source of behavioural data for ecosystem services analysis. Public geotagged photographs, for instance, represent a unique form of revealed preference data, available at wide scales and low cost. In Chapter 11, Sinclair et al. use online geotagged photographs from Flickr to evaluate and map recreation to the wetlands of Kerala state, India. Photographic metadata are used to: (1) estimate the most-visited wetlands; (2) uncover patterns of recreational use across different wetlands types; (3) investigate the relationship between visitation and various wetland and site-specific attributes such as area and water quality, through regression analysis; and (4) test scenarios of wetland improvement and their potential to impact upon visitation to five coastal wetlands presently suffering from poor water quality.

The models and methods used by scientists for environmental assessments and scenarios often face practical hurdles of application. Due to institutional barriers, including limited capacity, institutional silos and inaccessibility of high-tech quantitative modelling tools, advanced scenario planning methods may remain as academic exercises with little or no guidance for municipal planning agencies. Chapter 12 by Caughman et al. offers a framework for integrating assessments and scenarios into the municipal workflow to deal with grand environmental challenges. Their approach, the Scenario Collaboratory, works by engaging with a wide range of decision-makers and using academic knowledge to advance city planning outcomes. In this study, the Scenario Collaboratory framework was used to design and implement inter-agency scenario planning workshops in the City of Portland, which focused on resilience to and recovery from two major natural disasters: a magnitude 8.5 earthquake, and a 500-year (recurrence interval) flooding event. Through this in-depth case study, the authors draw out several principles about collaborative management and implementation that can support practical applications of scenario planning outcomes.

Chapter 13 by Ravera et al. reflects on the lessons learned from an iterative participatory modelling process applied to socio-environmental management in a semi-arid protected area in northern Nicaragua. The study was to understand vulnerability and support local stakeholders in adapting to socio-economic, environmental and policy pressures. The

chapter reports on the genesis and the approach developed. Researchers collaboratively co-created with stakeholders a set of conceptual models that identify key issues and make explicit local perceptions on the functioning of a complex social-ecological landscape. The authors coupled computer models to explore the implications of farmers' choices of land use and management expressed in a range of ecological and socio-economic indicators. Through 'playing with modelling' workshops they communicated back the results of simulations to discuss the current and potential future states of the local silvopastoral system. The authors reflect on the relevance of social learning processes and substantive outcomes in contexts with always incomplete knowledge, and discuss the challenge of bridging different knowledge systems together in order to assist decision-making, promote local interactions, and strengthen local agency and ownership of learning processes.

REFERENCES

- ADB (2003), *Environmental Assessment Guidelines*, Asian Development Bank, Manila, Philippines. <https://www.adb.org/sites/default/files/institutional-document/32635/files/environmental-assessment-guidelines.pdf>. Accessed on 24 October 2019.
- Arrhenius, O. (1921), 'Species and area', *Journal of Ecology*, **9** (1), 95–99.
- Barfield, M., R.D. Holt and R. Gomulkiewicz (2011), 'Evolution in stage-structured populations', *American Naturalist*, **177** (4), 397–409.
- Barton, N.H. (2001), 'The role of hybridization in evolution', *Molecular Ecology*, **10** (3), 551–568.
- Bateman, I., G.M. Mace, C. Fezzi, et al. (2014), 'Economic analysis for ecosystem assessments', in K.N. Ninan (ed.), *Valuing Ecosystem Services: Methodological Issues and Case Studies*, Cheltenham, UK and Northampton, MA, USA: Edward Elgar Publishing, pp. 23–77.
- Bellard, C., C. Bertelsmeier, P. Leadley, et al. (2012), 'Impacts of climate change on the future of biodiversity', *Ecology Letters*, **15** (4), 365–377.
- Botkin, D.B. (1993), *Forest Dynamics: An Ecological Model*, Oxford: Oxford University Press.
- Brotons, L., V. Christensen, N.H. Ravindranath, et al. (2016), 'Modelling impacts of drivers on biodiversity and ecosystem services', Chapter 4 in IPBES, *The Methodological Assessment Report on Scenarios and Models of Biodiversity and Ecosystem Services*, Bonn: Secretariat of the Intergovernmental Science–Policy Platform on Biodiversity and Ecosystem Services, United Nations, pp. 121–157.
- CBD (2019), 'What is impact assessment', Convention on Biological Diversity, UNEP, Montreal. <https://www.cbd.int/impact/whatis.shtml>. Accessed on 22 October 2019.
- Cheung, W.W.L., V.M.Y. Lam and D. Pauly (2008), *Dynamic Bioclimate Envelope Model to Predict Climate-induced Changes in Distributions of Marine Fishes and Invertebrates*, Vancouver: Fisheries Centre, University of British Columbia.

- Christensen, V. and C.J. Walters (2011), 'Progress in the use of ecosystem modelling for fisheries management', in V. Christensen and J.L. Maclean (eds), *Ecosystem Approaches to Fisheries: A Global Perspective*, Cambridge: Cambridge University Press, pp. 189–205.
- Christensen, V., C.J. Walters and D. Pauly (2005), *Ecopath with Ecosim: A Users Guide*, Vancouver: Fisheries Centre, University of British Columbia.
- Cramer, W., A. Bondeau, F.I. Woodward, et al. (2001), 'Global response of terrestrial ecosystem structure and function to CO₂ and climate change: results from the dynamic global vegetation models', *Global Change Biology*, **7** (4), 357–373.
- de Souza Muñoz, M.E., R. De Giovanni, M.F. de Siquiera, et al. (2011), 'open-Modeller: a generic approach to species potential distribution modelling', *GeoInformatica*, **15** (1), 111–135.
- Desrochers, R.E. and J.T. Kerr (2011), 'How and how much natural cover loss increases species richness', *Global Ecology and Biogeography*, **20** (6), 857–867.
- Duplisea, D.E., S. Jennings, K.J. Warr and T.A. Dinmore (2002), 'A size-based model of the impacts of bottom trawling on benthic community structure', *Canadian Journal of Fisheries and Aquatic Sciences*, **59** (11), 1785–1795.
- EEA (2009), *Looking Back on Looking Forward: A Review of Evaluative Scenario Literature*, EEA Technical Report No. 3/2009, Copenhagen: European Environment Agency.
- Elith, J. and J.R. Leathwick (2009), 'Species distribution models: ecological explanation and prediction across space and time', *Annual Review of Ecology, Evolution, and Systematics*, **40**, 677–697.
- Fernandes, P.G. and R.M. Cook (2013), 'Reversal of fish stock decline in the northeast Atlantic', *Current Biology*, **23** (15), 1432–1437.
- Ferrier, S. and A. Guisan (2006), 'Spatial modelling of biodiversity at the community level', *Journal of Applied Ecology*, **43** (3), 393–404.
- Gallego-Sala, A.V., J.M. Clark, J.I. House, et al. (2010), 'Bioclimatic envelope model of climate change impacts on blanket peatland distribution in Great Britain', *Climate Research*, **45** (1), 151–162.
- Getz, W.M. (1988), 'Harvesting discrete nonlinear age and stage structured populations', *Journal of Optimization Theory and Applications*, **57** (1), 69–83.
- Hanson, C., J. Ranganathan, C. Iceland and J. Finisdore (2012), *The Corporate Ecosystem Services Review: Guidelines for Identifying Business Risks and Opportunities arising from Ecosystem Change (version 2.0)*. <http://www.wri.org/publication/corporate-ecosystem-services-review>.
- Harfoot, M.B.J., D.P. Tittensor, T. Newboldt, et al. (2014), 'Integrated assessment models for ecologists: the present and the future', *Global Ecology and Biogeography*, **23** (2), 124–143.
- Harremoes, P. and R.K. Turner (2001), 'Methods for integrated assessment', *Regional Environmental Change*, **2** (2), 57–65.
- Hilborn, R. and C.J. Walters (1992), *Quantitative Fisheries Stock Assessment: Choice, Dynamics & Uncertainty*, Dordrecht: Chapman & Hall.
- Huth, N. and H.P. Possingham (2011), 'Basic ecological theory can inform habitat restoration for woodland birds', *Journal of Applied Ecology*, **48** (2), 293–300.
- Iglesia, M.N., J.A. Collazo and A.J. McKerrow (2012), 'Use of occupancy models to evaluate expert knowledge-based species habitat', *Avian Conservation and Ecology*, **7** (2), 5.
- IPBES (2016), *The Methodological Assessment Report on Scenarios and Models of Biodiversity and Ecosystem Services*, S. Ferrier, K.N. Ninan, P. Leadley,

- R. Alkemade et al., (eds), Secretariat of the Intergovernmental Science–Policy Platform on Biodiversity and Ecosystem Services, United Nations, Bonn.
- IPCC (2001), *The Third Assessment Report*, Geneva: IPCC.
- IPCC (2014), *Working Group II. Climate Change 2014: Impacts, Adaptation, and Vulnerability*, V.R. Barros, C.B. Field, D.J. Dokken, et al. (eds), Intergovernmental Panel on Climate Change (IPCC), Switzerland.
- Janse, J.J., J.J. Kuiper, M.J. Weijters, et al., (2015), ‘GLOBIO-Aquatic: a global model of human impact on the biodiversity of inland aquatic ecosystems’, *Environmental Science & Policy*, **48**, 99–114.
- Kok, M.T.J., K. Kok, G.D. Peterson, et al. (2017), ‘Biodiversity and ecosystem services require IPBES to take novel approaches to scenarios’, *Sustainability Science*, **12**, 177–181.
- Kooijman, S.L.A.M. (2009), *Dynamic Energy Budget Theory for Metabolic Organisation*, Cambridge: Cambridge University Press.
- Mouton, A.M., B. De Baets and P.L.M. Goethals (2009), ‘Knowledge-based versus data-based fuzzy habitat suitability models for river management’, *Environmental Modelling & Software*, **24** (8), 982–993.
- Neugarten, R.A., P.F. Langhammer, E. Osipova, et al. (2018), *IUCN Report on Tools and Methods for Evaluating Ecosystem Services*, Best Practice Protected Area Guideline Series No. 28, Gland, Switzerland: International Union for Conservation of Nature and Natural Resources. <https://portals.iucn.org/library/sites/library/files/documents/PAG-028-En.pdf>.
- Notaro, M., A. Mauss and J.W. Williams (2012), ‘Projected vegetation changes for the American southwest: combined dynamic modelling and bioclimate envelope approach’, *Ecological Applications*, **22** (4), 1365–1388.
- Parrott, L. (2011), ‘Hybrid modelling of complex ecological systems for decision support: recent successes and future perspectives’, *Ecological Informatics*, **6** (1), 44–49.
- Pearce, D., G. Atkinson and S. Mourato (2006), *Cost–Benefit Analysis and the Environment – Recent Developments*, Paris: Organisation for Economic Co-operation and Development (OECD).
- Pearson, R.G. and T.P. Dawson (2003), ‘Predicting the impacts of climate change on the distribution of species: are bioclimate envelope models useful?’, *Global Ecology and Biogeography*, **12** (5), 361–371.
- Peh, K.S.H., A.P. Balmford, R.B. Bradbury, et al. (2014), *Toolkit for Ecosystem Service Site-Based Assessment (TESSA)*, version 1.2, Cambridge.
- Periera, H.M., P. Leadley, V. Proenca, et al. (2010), ‘Scenarios for global biodiversity in the 21st century’, *Science*, **330** (6010), 1496–1501.
- Pereira, H.M., S. Ferrier, M. Walters, et al. (2013), ‘Essential biodiversity variables’, *Science*, **339** (6117), 277–278.
- Peterson, A.T. (2006), ‘Uses and requirements of ecological niche models and related distributional models’, *Biodiversity Informatics*, **3**, 59–72.
- Peterson, G., J. Kabubo-Mariara, N. Crossman, et al. (2016), ‘Modelling consequences of change in biodiversity and ecosystems for nature’s benefits to people’, Chapter 5 in IPBES, *The Methodological Assessment Report on Scenarios and Models of Biodiversity and Ecosystem Services*, Bonn: Secretariat of the Intergovernmental Science–Policy Platform on Biodiversity and Ecosystem Services, United Nations, pp. 161–193.
- Phillips, S.J., R.P. Andersen and R.E. Schapire (2006), ‘Maximum entropy modelling of species geographic distributions’, *Ecological Modelling*, **190** (3–4), 231–259.

- Polechová, J. and N.H. Barton (2015), 'Limits to adaptation along environmental gradients', *Proceedings of the National Academy of Sciences*, 6401–6406. doi:10.1073/pnas.1421515112.
- Priess, J.A. and J. Hauck (2014), 'Integrative scenario development', *Ecology and Society*, **19**, 14.
- Rochet, M.J., J.S. Collie, J. Simon and S.J. Hall (2011), 'Does selective fishing conserve community biodiversity? Predictions from a length-based multi species model', *Canadian Journal of Fisheries and Aquatic Sciences*, **68** (3), 469–486.
- Rosa, I.M.D., H.M. Periera, S. Ferrier, et al. (2017), 'Multiscale scenarios for nature futures', *Nature Ecology & Evolution*, **1**, 1416–1419.
- Rosenzweig, M.L. (2010), *Species Diversity in Space and Time*, Cambridge: Cambridge University Press.
- Sharp, R., H.T. Tallis, T. Ricketts, et al. (2018), *INVEST 3.6.0 User's Guide*, The Natural Capital Project, Stanford University, University of Minnesota, The Nature Conservancy, and World Wildlife Fund.
- Sinclair, A.R.E. and J.M. Gosline (1997), 'Solar activity and mammal cycles in the northern hemisphere', *American Naturalist*, **149** (4), 776–784.
- Spangenberg, J.H., A. Bondeau, T.H. Carter, et al. (2012), 'Scenarios for investigating risks to biodiversity', *Global Ecology and Biogeography*, **21**, 5–18.
- Stehfest, E., D. van Vuuren, T. Kram, et al. (2014), *Integrated Assessment of Global Environmental Change with IMAGE 3.0. Model Description and Policy Applications*, The Hague: PBL, Netherlands Environmental Assessment Agency.
- Stockwell, D. and D. Peters (1999), 'The GARP modelling system: problems and solutions to automated spatial prediction', *International Journal of Geographical Information Science*, **13** (2), 13–158.
- Tilman, D., J. Fargione, B. Wolff, et al. (2001), 'Forecasting agriculturally driven global environmental change', *Science*, **292** (5515), 281–284.
- UNEP (2019), *Guidelines for Conducting Integrated Environmental Assessments*, Nairobi: United Nations Environment Programme (UNEP).
- van der Heijden, K., R. Bradfield, G. Burt, et al. (2002), *The Sixth Sense: Accelerating Organisational Learning with Scenarios*, Chichester: Wiley & Sons.
- van Notten, P. (2005), 'Scenario development: a typology of approaches'. Available at <https://www1.oecd.org/Site/Schoolingfortomorrowknowledgebase/futuresthinking/Scenarios/3724643.pdf>. Accessed on 1 November 2019.
- van Vuuren, D.P., O.E. Sala and H.M. Periera (2006), 'The future of vascular plant diversity under four global scenarios', *Ecology and Society*, **11** (2), 25. <http://www.ecologyandsociety.org/vol11/iss2/art25/>.
- van Vuuren, D.P., M.T.J. Kok, B. Girod, et al. (2012), 'Scenarios in global environmental assessments: key characteristics and lessons for future use', *Global Environmental Change*, **22**, 884–895.
- Villa, F., K.J. Bagstad, B. Voigt, et al. (2014), 'A methodology for adaptable and robust ecosystem services assessment', *PLoS ONE*, **9** (3), e91001. doi: 10.1371/journal.pone.0091001.
- <https://seea.un.org/>, 'What is SEEA'. Accessed on 24 October 2019.
- <https://gnhusa.org/genuine-progress-indicator/>, 'Genuine Progress Indicator'. Accessed on 24 October 2019.
- <https://www.nature.com/subjects/ecological-modelling>. Accessed on 16 November 2019.